



Original Article

MUTAGENIC EFFECTS OF FAST NEUTRON IRRADIATION ON SOME GROWTH AND YIELD PARAMETERS AT M₁ and M₂ OF LIMA BEAN (*Phaseolus lunatus* Linn.)

***¹Ikani, V. O., ²Adelanwa, M. A. and ²Aliyu, R. E.**

¹Department of Biology, Ahmadu Bello University, Zaria, Nigeria

²Department of Botany, Ahmadu Bello University, Zaria, Nigeria

Submitted: February 2017; Accepted: April 2017; Published: June, 2017

ABSTRACT

The studies of induced mutation in *Phaseolus lunatus* L. was conducted at the botanical garden, Department of Botany, Ahmadu Bello University, Zaria. Dry healthy seeds of *P. lunatus* obtained from local farmers with in Zaria were exposed to Fast Neutron irradiation at different doses (0.00Sv, 0.16Sv, 0.32Sv, 0.49Sv and 0.65Sv). The irradiated seeds were sown in 150 polythene bags which were half filled with soil and arranged using a Randomized Complete Block design. The result of this studies showed both positive and negative shift in agronomic characters with plants exposed to lower doses (0.16Sv and 0.32Sv) showing better performance in most agronomic trait in both mutagenic generations while higher dose (0.65Sv) recorded the least performance in most agronomic traits. Chlorophyll mutants such as Zonata, Xantha and Maculata as well as Morphological mutants which includes invaginated leaf margins, bifurcated leaf apex and bifoilate leaves were observed. However, lower doses (0.16Sv and 0.32Sv) of the mutagen were more favourable in inducing beneficial mutation on the growth and yield trait of *P. lunatus*. The effect of the mutagen was significant at $p \leq 0.05$ in plant height 4 weeks after sowing, days to first flowering, days to 50% flowering, pod length, pod diameter, pod weight and 100 seeds weight.

Keywords: Fast neutron, *P. lunatus*, Seeds, Agronomic traits, Chlorophyll mutants and Morphological mutants.

***Corresponding author:** vonuche@gmail.com

INTRODUCTION

Phaseolus lunatus is commonly called lima bean, butter bean, sieva bean, sugar bean, Madagascar bean and java bean in

English, Wake Rumpa in Hausa, Ukpa in Igbo (Ikechukwu, 2010), Papala in Yoruba (Adegbehingbe, 2013). It belongs to the family Fabaceae (USDA, 2012). Like other

grain legumes, lima beans are relatively rich in protein. They contain about 25% protein in the dry matter, a value comparable to that of peas (*Pisum sativum*) and cowpeas (*Vigna unguiculata*), dried vines may be used as animal fodder (Azeke *et al.*, 2011). Immature sprouts, leaves, and pods are consumed in Asia with young leaves and pods steamed and green shelled beans cooked as a vegetable. Lima bean helps restore soil fertility by shedding its many leaves that decay and enrich the soil. It is also used as a green manure and cover crop to prevent soil erosion from heavy rainfall in rubber plantations Oliveira *et al.* (2004).

Mutation breeding is a proven supplement and an effective substitute of conventional breeding so as to confer specific improvement in a variety without significantly affecting its acceptable phenotype (Micke, 1998). Physical mutagens like x-rays, gamma rays, fast neutrons, thermal neutrons, ultraviolet and beta radiations have been frequently used to induce mutation (Yakoob and Rashid, 2001). Fast neutrons are produced by cyclotrons or atomic reactors as a result of radioactive decay of heavier elements. Neutrons are uncharged particles, and are highly penetrating in biological tissues; fast and thermal neutrons are densely ionizing radiations (Singh, 2013). The state of nutrition in Nigeria is still characterized by inadequate calorie and protein supplies (FAO, 2004). Despite the great potential of lima bean, it is highly underutilized in Nigeria and it has not received much attention in terms of crop improvement; local cultivars are still being grown by the farmer and so yield is low (Akande and Balogun, 2007). In an

attempt to widen the narrow genetic base, food and agricultural scientists are screening lesser known and under-exploited native plants for possible potential sources of food (Ezeagu and Ologhobo, 1995; Vietmeyer and Janick, 1996; Murray *et al.*, 2001). One of such crop identified in this context in Nigeria is lima bean, though a minor crop, has been an important source of plant protein to millions of Nigerians (Ikechukwu *et al.*, 2010). Thus, this research was targeted towards determining the mutagenic effects of fast neutrons on some growth and yield parameters of *P. lunatus*

MATERIALS AND METHODS

Seed Source and Mutagenic Treatment

Healthy landraces of *P. lunatus* were obtained from local farmers in Zaria, Kaduna State, Nigeria and landrace identity was confirmed at International Centre for Tropical Agriculture (CIAT), Cali, Colombia. About 0.25kg of healthy dry seeds of *P. lunatus* was exposed to Fast neutrons from Americium Beryllium source ($^{241}\text{Am}/\text{Be}$), with a flux of $1.5 \times 10^4 \text{ncm}^{-2}\text{s}^{-1}$ at Center for Energy Research and Training (CERT), Ahmadu Bello University, Zaria for 0hrs, 2hrs, 4hrs, 6hrs and 8hrs equivalent to 0.00Sv, 0.16Sv, 0.32Sv, 0.49Sv and 0.65Sv doses respectively

Experimental design and sowing of seeds

Top soil was collected from uncultivated land within the botanical garden in Ahmadu Bello University, Zaria and after treatment it was filled into one hundred and fifty (150) polythene bags (51.5 X 38.3cm) arranged in a randomized complete block design (RCBD) with three replications for each treatment to raise M₁ and M₂ generation. The polythene bags were half filled each weighing 8.5kg and placed at a spacing of 10 x 90 cm. A total of 1.35 x 10⁻³ hectares (13.5msq) of land were used for this research. Three seeds were sown in each bag which was later thinned to one per bag.

Sampling Collection

Agro-morphological data such as Percentage germination, plant height 4 weeks after sowing (WAS), 8WAS, 12WAS, number of days to first flowering, number of days to 50% flowering, number of pods per plant, pod length, pod weight, pod diameter, number of seeds per pod and 100 seeds weight.

Data Analyses

Agro-morphological data obtained was subjected to one way analysis of variance (ANOVA) using SAS (2002) version 9.1 to determine the significant effect of different concentrations of the mutagens ($p \leq 0.05$) and Duncan's Multiple Range test (DMRT) was used to separate the means obtained, where significant.

RESULTS

Effects of Fast Neutron on some growth and yield parameters of *P. lunatus* at M₁Generation

The result of the effect of different doses of fast neutron irradiation on the agronomic traits of *P. lunatus* is shown in

Table 1. Plant height at 8 and 12 WAS, days to first flowering, days to 50% flowering, number of pods per plant, pod length and pod diameter were significantly different ($P \leq 0.05$). The highest plant height (157.32cm) and least plant height (76.25cm) at 8WAS were observed at 0.32Sv and the control respectively. Plant height 12WAS increased with an increase in dose, 0.32Sv recorded the highest plant height (179.79cm) while the control recorded the least height (105cm). However, the highest number of days to first (86.00) and 50% (93.00) flowering was recorded at 0.49Sv dose while the fewest (73.33 and 80.67) respectively was at 0.16Sv. The number of pods per plant decreased with an increase in dose of the mutagen, where the control produced the highest (17.80) and the least (10.40) was produced at 0.65Sv. Pod length also varied with dose although not linearly, the highest pod length (6.59cm) was recorded at 0.49Sv and the least by the control (5.36cm). On the other hand 0.49Sv and the control recorded the highest (1.77cm) and least (1.43cm) pod diameter respectively. Pod weight was observed to have increased with an increase in dose of fast neutrons; 0.49Sv recorded the highest mean value 7.20g) while the least mean value (5.50) was recorded by the control treatment.

Effects of Fast Neutron on some growth and yield parameters of *P. lunatus* at M₂Generation

Number of days to first flowering, days to 50% flowering, pod length, pod diameter, pod weight and 100 seed weight showed significant difference ($P \leq 0.05$) (Table 2) The control treatment showed the highest number of days to first (54.33) and 50% flowering (63.67) while the fewest

number of days to first flowering (50.00) was recorded at 0.16Sv and 0.65Sv and 0.49Sv recorded the fewest number of days to 50% flowering (59.00). Although, not linearly the pod length increased with an increase in dosage; 0.32Sv produced longest pod (5.79cm) and the shortest pod (4.73) was produced by the control treatment. Similarly, 0.32Sv produced the pods with the widest diameter (1.74)

while the control had the narrowest diameter (1.58cm), this was followed closely by 0.16Sv with a diameter of 1.60cm. The highest pod weight (6.47g) was observed at 0.33Sv and the least (4.27g) in the control treatment, 100 seed weight also varied amongst the different doses of fast neutrons with 0.16Sv weighing the highest (32.33g) and 0.65Sv weighing the least (28.33g)

Table 1: Effects Fast Neutrons on Agronomic Traits of *Phaseolus lunatus* at M₁ Generation

TRT	PGERM (%)	PH4 (cm)	PH8 (cm)	PH12 (cm)	DFF	D50%F	NPPP	PDLT (cm)	PDDIA (cm)	PDWT (g)	NSPP	100SWT (g)
0.0Sv	86.67 ^a	32.23 ^a	76.25 ^b	105.00 ^b	77.67 ^b	85.00 ^b	17.80 ^a	5.36 ^c	1.43 ^d	5.50 ^b	2.60 ^a	30.00 ^a
0.16Sv	83.33 ^a	40.80 ^a	85.18 ^b	151.85 ^a	73.33 ^c	80.67 ^c	14.40 ^b	5.97 ^b	1.60 ^{bc}	6.20 ^{ab}	2.87 ^a	30.67 ^a
0.32Sv	80.00 ^a	28.47 ^a	157.32 ^a	179.97 ^a	79.67 ^b	86.67 ^b	13.13 ^{bc}	6.08 ^b	1.69 ^{ab}	6.77 ^a	2.80 ^a	31.00 ^a
0.49Sv	80.00 ^a	31.97 ^a	113.25 ^{ab}	169.74 ^a	86.00 ^a	93.00 ^a	10.53 ^c	6.59 ^a	1.77 ^a	7.20 ^a	2.80 ^a	32.33 ^a
0.65Sv	80.00 ^a	29.50 ^a	103.80 ^b	158.13 ^a	79.67 ^b	87.00 ^b	10.40 ^c	5.69 ^{bc}	1.55 ^{cd}	5.60 ^b	2.67 ^a	32.33 ^a
SEM±	7.89	4.73	15.18	13.38	1.01	1.10	0.92	0.13	0.04	0.34	0.25	1.19

NOTE: Means with the same letter within a column are not significantly different at $P \leq 0.05$

TRT- treatment, PGERM- percentage germination, PH4-plant height at 4 weeks, PH8-plant height at 8 weeks, PH12-plant height at 12weeks, DFF-days to first flowering, D50%F-days to 50% flowering, NPPP-number of pods per plant ,PDLGTH-pod length, PDDIA- pod diameter, PDWT-pod weight, NSPP-number of seeds per pod 100SWT-one hundred seeds weight

Table 2: Effects Fast ``Neutrons on Agronomic Traits of *Phaseolus lunatus* at M₂ Generation

TRT	PGERM (%)	PH4 (cm)	PH8 (cm)	PH12 (cm)	DFF	D50%F	NPPP	PDLT (cm)	PDDIA (cm)	PDWT (g)	NSPP	100SWT (g)
0.0Sv	93.33 ^a	37.93 ^b	98.88 ^a	112.63 ^a	54.33 ^a	63.67 ^a	6.87 ^a	4.73 ^c	1.58 ^b	4.27 ^c	2.20 ^a	29.90 ^b
0.16Sv	100.00 ^a	60.66 ^a	100.35 ^a	118.26 ^a	50.00 ^c	61.33 ^b	6.60 ^a	5.34 ^b	1.60 ^b	4.60 ^b	2.53 ^a	32.33 ^a
0.32Sv	93.33 ^a	50.01 ^{ab}	103.74 ^a	117.30 ^a	50.33 ^c	62.33 ^{ab}	7.47 ^a	5.79 ^a	1.74 ^a	6.47 ^a	2.67 ^a	30.33 ^{ab}
0.49Sv	90.00 ^a	57.32 ^{ab}	94.94 ^a	111.87 ^a	51.67 ^b	59.00 ^c	8.13 ^a	5.40 ^b	1.68 ^a	6.20 ^{ab}	2.67 ^a	29.67 ^b
0.65Sv	96.67 ^a	51.45 ^{ab}	99.80 ^a	109.93 ^a	50.00 ^c	59.67 ^c	8.07 ^a	5.61 ^{ab}	1.68 ^a	6.23 ^{ab}	2.60 ^a	28.33 ^b
SEM±	3.80	6.33	4.84	7.15	0.38	0.46	1.09	0.08	0.02	0.24	0.17	0.68

NOTE: Means with the same letter within a column are not significantly different at $P \leq 0.05$

TRT- treatment, PGERM- percentage germination, PH4-plant height at 4 weeks, PH8-plant height at 8 weeks, PH12-plant height at 12weeks, DFF-days to first flowering, D50%F-days to 50% flowering, NPPP-number of pods per plant ,PDLGTH-pod length, PDDIA- pod diameter, PDWT-pod weight, NSPP-number of seeds per pod 100SWT-one hundred seeds weight

Chlorophyll mutant such as Zonata showed dark light coloured bands like distinct zones spread over the leaf, Xantha showed light yellow colour indicating that

the carotenoids prevailed over the chlorophyll, Maculata mutant showed chlorophyll or carotene destruction in the form of dots on the surface of the leaves.

However, morphological mutants such as bifoilate leaves, invaginated leaf margin,

double invaginated leaf margin and bifurcated leaf apex were observed.



Plate 1: Control

Plate 2: Zonata

Plate 3: Xantha

Plate 4: Maculata



Plate 5: Double
invaginated margin

Plate 6: Bi-furcated
apex

Plate 7: invaginated
margin

Plate 8: Bifoilate leaf

DISCUSSION

The increase in plant height at 8WAS, 12WAS, pod length, pod diameter and pod weight with increase in exposure to fast neutrons irradiation at M_1 generation, may be due to the alteration of their genome integrated by environmental signals as reported by Uno *et al.* (2001); probably by increasing the rate of cellular division and expansion at their meristematic regions. This is also in agreement with the findings of Hoballah (1999) who reported increase in plant height of Sesame due to radiation

mutagenesis, Lysenkov (1989) who induced a wide range of viable mutants and observed increase in wheat height, Adamu *et al.* (2002) when they irradiated the seeds of five varieties of groundnut (*Arachis hypogaea*) with gamma rays Cobalt 60 (^{60}Co) source, Falusi *et al.* (2012) when they irradiated the seeds of *Capsicum annum* with fast neutrons but disagrees with the findings of Adamu (2004) who exposed (pop-corn maize) to thermal neutron and gamma irradiation.

Days to first flowering and 50% flowering which also increased with an increase in

exposure to irradiation might be due to inhibitory effects of the mutants and physiological changes in the plant as a result of delayed cellular division prior to flowering and reproductive stage especially at higher doses, this conforms to the findings of Sadiq (2015) and contradicts the findings of Mensah *et al.* (2007), in sesame and Mathew (2014), in pigeon pea. The number of pods per plant when compared to the control decreased with an increase in dosage of the radiation, this might be as a result of increased injuriousness on exposure to higher doses of fast neutrons irradiation, this is similar to the findings of Muhammad *et al.* (2013) when they treated *Sesamum indicum* with fast neutrons irradiation and observed a decrease in the number of capsules per plant with an increase in dose and disagrees with the findings of Mathew (2014), in pigeon pea and Sadiq (2015), in soya bean.

Percentage germination, Plant height 8WAS, 12WAS, Number of pods per plant and number of seeds per pod at M₂ generation were not significantly different ($P > 0.05$) this might be as a result of the fact that the dosage of fast neutrons was not adequate enough to cause reasonable level of variability in this character. This is similar to the findings of Abdullahi *et al.* (2015) who treated *Phaseolus vulgaris* with gamma rays. Days to first flowering and 50% flowering were significantly different when compared to the control and this maybe as a result of the heritable physiological changes that must have occurred at the M₁ generation. Pod length, Pod weight, Pod diameter and 100 seed weight were significantly different from the control and this study agrees with the

findings of Elangovan and Pavadai (2015).

Chlorophyll and morphological mutants were classified in accordance with the system of Gustafsson (1940 and 1941). Macro mutations are generally used to evaluate the genetic effects of various mutagens. Gaul (1964) reported that chlorophyll mutations are employed as markers for the evaluation of gene action of mutagenic factors in inducing mutation. Chlorophyll mutations provide one of the most dependable indices for the evaluation of genetic effects of mutagenic treatment and have been reported in various pulse crops by several workers including Gautam *et al.* (1992). They do not have any major implication in mutation breeding but is an indication of the effectiveness and efficiency of the mutagen in inducing mutation.

CONCLUSION

This research concluded that the effects of fast neutrons irradiation varied with the various doses. Lower doses (0.16Sv and 0.32Sv) of the mutagen were more favourable in inducing beneficial mutation on the growth and yield trait of *P. lunatus*. The effect of the mutagen was significant at $p \leq 0.05$ in plant height at 4 weeks, days to first flowering, days to 50% flowering, pod length, pod diameter, pod weight and 100 seeds weight at the M₂ generation.

ACKNOWLEDGEMENTS

We are grateful to the Departments of Biology and Botany Ahmadu Bello University, Zaria, Nigeria for providing the necessary laboratory facilities and land for the field experiment.

REFERENCES

- Abdullahi, S., Abubakar, B. Y., Adelanwa, M. A., Shehu, S. A., Zangoma, I. M. and Amshi, A. M. (2015). Effects of gamma rays and Sodium azide on yield parameters of *Phaseolus vulgaris* L. *International Journal of Research*, 2(11): 702-714.
- Adamu, A. K. (2004). Gamma rays and thermal neutron induced mutants in popcorn (*Zea mays* var. *praecox* Sturt). *Nigerian Journal of Scientific Research*, 4(2): 52-63.
- Adamu, A. K., Olorunju, P. E., Bate, J. A. and Ogunlade, O. T. E. (2002). Radio sensitivity and effective dose determination in Groundnut (*Arachis hypogaea* L.) Irradiated with Gamma Rays (Cobalt 60) sources. *Journal of Agricultural and Environment*, 3(1): 71-84.
- Adegbekingbe, K. T. (2013). Microbial and Nutrient Studies of Fermented Cooked Lima Bean (*Phaseolus lunatus* L.) seeds. *Global Journal of Biology, Agriculture and Health Sciences*, 2(2): 94-101.
- Akande, S. R. and Balogun, M.O. (2007). Evaluation and heritability studies of local bean (*Phaseolus lunatus* L.) cultivars from south-west Nigeria. *Revista UDO Agricola*, 7(1): 22-28.
- Azeke, M. A., Elsanhoty, R. M., Egielewa, S. J. and Eigbogbo, M. U. (2011). The effect of germination on the phytase activity, phytate and total phosphorus contents of some Nigerian-grown grain legumes. *Journal of the Science of Food and Agriculture*, 91: 75-79.
- Elangovan, R. and Pavadai, P. (2015). Studies on induced chemical mutagenesis in Bhendi (*Abelmoschus esculentus* (L.) Moench). *International Journal of Modern Biology and Medicine*, 6(1): 30-37.
- Ezeagu, I. E. and Ologhobo, A. D. (1995). Proximate composition of unfamiliar plant seeds in Nigeria. A short report. *Poland Journal of Food and Nutrition Sciences*, 4(5): 79-83.
- Falusi, O. A., Daudu, O. A. Y. and da Silva, J. A. T. (2012). Effect of exposure time of fast neutron irradiation on growth and yield parameters of *Capsicum annum* and *Capsicum frutescens*. *African Journal of Plant Science*, 6(9): 251-255.
- Food Agricultural Organization (2004). The state of food insecurity in the World 2004, FAO, Rome, Italy.
- Gaul, H. (1964). Mutations in plant breeding. *Radiation Botany*, 4(3): 155-232.
- Gautam, A. S., Sood, K. C. and Richarria, A. K. (1992). Mutagenic effectiveness and efficiency of gamma rays, ethyl methane sulphonate and their synergistic effects in black gram (*Vigna mungo* L.). *Cytologia*, 57: 85-89.
- Gustafsson, Å. (1940). The mutation system of the chlorophyll apparatus. *Lunds Universitets Årsskrift, N. F. Avd. 2. Bd. 36*: 1 - 40.
- Gustafsson, Å. (1941). Mutation experiments in barley. *Hereditas*, 27: 225 - 242.

- Hoballah, A. A. (1999). Selection and Agronomic evaluation of Induced Mutant lines of Sesame. In. Induced Mutations for Sesame Improvement IAEA-TECDOC, IAEA, Vienna. Pp. 71-84.
- Ikechukwu, E. E. and Madu, D. I. (2010). Biochemical Composition and Nutritional Potential of Ukpa, A variety of Tropical Lima Bean (*Phaseolus lunatus*) from Nigeria - A Short Report. *Polish Journal of Food and Nutritional Science*, 60(2): 231-235.
- Lysenkov, V. I. (1989). Experimental production of mutants of winter breeds wheat, as breeding material. *Vavilour*; 187: 17-20.
- Mathew, B. A. (2014). Morphological and Cytogenetics studies on the effect of Sodium Azide and Gamma irradiation treatments on Pigeon pea (*Cajanus cajan* (L.) Millspaugh). Unpublished MSc. Thesis, Ahmadu Bello University, Zaria, Nigeria, 78pp.
- Mensah, J. K., Obadoni, B. O., Akomeah, P.A., Ikhajiagbe, B. and Ajibulu, J. (2007). The effects of sodium azide and Colchicine treatments on morphological and yield traits of sesame seed (*Sesamum indicum* L.). *African Journal of Biotechnology*, 6(5): 534-538.
- Micke, A. (1998). Genetic Improvement of grain legumes using induced mutation. An overview. In, mprovement of Grain Legume Production using Induced mutations, IAEA, Vienna, pp.1 - 51.
- Muhammad, L. M., Falusi, O. A., Daudu, O. A. Y., Gado, A. A., Lateef, A. A. and Yahaya, S. A. (2013). Radiation induced polygenic mutation in two common Nigerian sesame (*Sesamum indicum* L.) cultivars. *International Journal of Biotechnology and Food Science*, 1(2): 23-28.
- Murray, S. S., Schoeninger, M. J., Bunn, H. T., Pickering, T. R. and Marlett, J. A. (2001). Nutritional composition of some wild plant foods and honey used by Hadza foragers of Tanzania. *Journal of Food Composition and Analysis*, 14: 3-13.
- Oliveira, A. P., Alves, E. U., Alves, A. U., Dornelas, C. S. M., Silva, J.A., Porto, M.L. and Alves, A.V. (2004). Produção de feijão-fava em função do uso de doses de fósforo em um Neossolo Regolítico. *Horticultura Brasileira*, 22: 543-546.
- Sadiq, S. A. (2015). Effects of gamma radiation and colchicine on agronomic Performance of soy-bean [*Glycine max* (L.) Merrill] varieties. Unpublished M.Sc. thesis, Department of Biological Sciences, Ahmadu Bello University, Zaria, Nigeria, 67Pp.
- Singh, B. D. (2013). *Plant Breeding, Principles and Methods*. Kalyani Publishers New Delhi- 110 002, India. 923Pp.
- Statistical Analysis Software (SAS) (2002) version 9.1. SAS Institute, North Carolina State University.
- United States Department of Agriculture (USDA) (2012). Natural Resources Conservation Service. <https://plants.usda.gov/core/profile>
- Uno, G., Storey, R. and Moore, R. (2001). *Principles of Botany*, McGraw Hill New York Pp. 1- 550

Vietmeyer, N. and Janick, J. (1996). New crops, 1996. in, Proceedings of the Third National Symposium, Indianapolis, Indiana, USA, 22-25 October 1996, *American Society for Horticultural Science Alexandria, USA*, pp. 2-8.

Yakoob, M. and Rashid, A. (2001). Induced Mutation Studies in Some Mung bean (*Vigna radiata* L.) Wilczex Cultivars. *Online Journal of Biological Sciences*, 1: 805-808.