

SOME COMBINATORIAL RESULTS ON ALTERNATING SEMIGROUPS

BAKARE, G. N. & MAKANJUOLA, S. O.
 Department of Mathematics
 University of Ilorin
 E-mail: bakaregattanaimat@gmail.com,
somakanjuola@unilorin.edu.ng

Abstract

We constructed the elements of A_n^E alternating semigroup on n-objects. We also investigated the combinatorial properties of the idempotent and nilpotent elements.

Keywords: Semigroup, Symmetric Inverse semigroup, Alternating semigroup, Idempotent and Nilpotent Element.

Introduction

Let $X_n = \{1, 2, \dots, n\}$. Then a (partial) transformation $\alpha : \text{Dom}\alpha \subseteq X_n \rightarrow \text{Im}\alpha$ is said to be full or total if $\text{Dom}\alpha = X_n$, otherwise is called Strictly Partial. It is denoted by P_n , when it is partial, T_n when it is full or total and C_n when it is one to one partial transformation semigroups. It is well known fact that if $S = C_n$ then $|S| = C_n = \sum_{r=0}^n \binom{n}{r}^2 r!$ Borwein [1989].

Alternating group is the group formed by even permutation on an n-objects denoted by A_n . A permutation is said to be even if it can be expressed as a product of an even number of transpositions.

The idea of even permutation can be extended to that of an even chart. The even charts in C_n partial one to one forms the alternating semigroup on n-objects $A_n^E \subseteq C_n$. As expected $A_n^E \cap S_n = A_n$.

Even Chart: for $\alpha \in C_n$ and $m \in \text{dom}\alpha$, " α moves m" when $m\alpha \neq m$ and that " α fixes m" when $m\alpha = m$. For instance let $N = \{1, 2, \dots\}$ then the transposition $(i, j) \in S_n$ moves i and j while fixing each $k \in N - ij$

In other words, A chart is said to be even if it can be expressed as a product of an even number of transpositional.

Transposition: Transposition is a circuit of length two it formed by $\{i, j\}$.

Transpositional: A transpositional is a chart either transposition $\{i, j\}$ or semitransposition $\{i, j\}$.

Idempotent: An element b in a transformation semigroup S is called an idempotent ($b^2 = b$) if and only if $\text{Im}(b) = F(b)$

Nilpotent: an element X in a semigroup with zero S, is a nilpotent if their exists a positive integer k such that $X^k = 0$.

Borwein (1989a) derived the formular for the number of semigroup $|S|$ in partial one-one transformation to be

$$C_n = \sum_{r=0}^n \binom{n}{r}^2 r!$$

$$C_0 = 1, C_1 = 2$$

Borwein (1989b) also derived the formula for the number of semigroup of the order decreasing partial one-one transformation to be the bell number

Where

$$B_{n+1} = \sum_{r=0}^n \binom{n}{r} B_r B_0 = 1 = B_1$$

Borwein (1989c) also derived the formula for the number of Idempotent $|E(S)|$ of the order-decreasing partial one-one transformation to be 2^n for $n \geq 1$

Umar in (1992) derived the formula for the Nilpotent $|N(S)|$ of the order –decreasing partial one-one transformation to be bell number

$$B_n$$

Umar (2004) reported that the formula for the number of Idempotent $|E(S)|$ in partial one-one transformation to be 2^n for $n \geq 1$

Laradji (2007) also derived the formula for the number of nilpotent element for $|N(S)|$ in partial one-one transformation to be

$$|NC_n| = \sum_{r=0}^{n-1} \binom{n}{r} \binom{n-1}{r} r! = u_n$$

Where

$$U_n = (2n-1)U_{n-1} - (n-1)(n-2)U_{n-2} U_0 = 1, U_1 = 1$$

Garba (1994a) derived the formula for the number of semigroup of the order preserving partial one-one transformation to be

$$CO_n = \binom{2n}{n}$$

Garba (1994b) also derived the formula for the number of Idempotent $|E(S)|$ for the order preserving partial one-one transformation to be 2^n for $n \geq 1$.

Ganyuushkin in (2003a) derived the formula for the number of semigroup of the order preserving and order decreasing partial one-one transformation to be I_{n+1} where

$$I_n = \frac{1}{n} \binom{2n}{n-1}$$

Ganyuushkin in (2003b) also derived the formula for the number of Idempotent $|E(S)|$ of the order preserving and order decreasing partial one-one transformation to be 2^n for $n \geq 1$

Stephen lmscomb derived the formula for the number of alter=natingsemigroup in partial one-one transformation to be

$$|A_n^e| = \frac{n!}{2} + \frac{n^2}{2} (n-1)! + \sum_{r=0}^{n-2} \binom{n}{r}^2 r!$$

Identification of Elements of A_n^e in C_n

We construct the elements of partial one-one transformation semigroup C_n and identifies elements of alternating semigroup (+) in it. The idempotent and nilpotent elements in A_n^e are also identified and marked (-) and (*) respectively.

The List

The elements in C_1

For $n=1$, has 2 elements.

C_1	{1}
	$\binom{1}{1} \binom{1^+}{1}$

The elements in C_2

For $n=2$, has 7 elements.

C_2	{1, 2}
	$\binom{12^+}{12} \binom{12}{21} \binom{12^+}{1-} \binom{12}{-1} \binom{12}{2-} \binom{12^+}{-2} \binom{12^+}{--}$

The elements in C_3

For $n=3$, has 34 elements

C_3	{1, 2, 3}
	$\binom{123}{123^+} \binom{123}{132} \binom{123}{213} \binom{123^+}{231} \binom{123^+}{312} \binom{123}{321} \binom{123^+}{12-} \binom{123}{1-2}$
	$\binom{123^+}{-12} \binom{123}{13-} \binom{123^+}{1-3} \binom{123}{-13} \binom{123}{21-} \binom{123^+}{21-} \binom{123}{-21} \binom{123^+}{23-}$
	$\binom{123}{2-3} \binom{123^+}{-23} \binom{123^+}{31-} \binom{123}{3-1} \binom{123^+}{-31} \binom{123}{32-} \binom{123}{3-2} \binom{123^+}{-32}$
	$\binom{123^+}{1--} \binom{123^+}{-1-} \binom{123^+}{--1} \binom{123^+}{2--} \binom{123^+}{-2-} \binom{123^+}{--2} \binom{123^+}{3--}$
	$\binom{123^+}{-3-} \binom{123^+}{--3} \binom{123^+}{---}$

The elements of partial one-one transformation semigroup of C_4 and C_5 were also generated but because of the space we could not display it. The table below shown the results

The Elements of Partial one-one Transformation Semigroup and Alternating Semigroup

Table 1

N	C_n	A_n^c
1	2	1
2	7	4
3	34	22
4	209	149
5	1546	1186

From the table 1 above, the sequences 2, 7, 34, 209, 1546,... the general formular which was derived by [1] was

$$C_n = \sum_{r=0}^n \binom{n}{r}^2 r!$$

$n \geq 2$, moreover, for

$$C_n = 2nC_{n-1} - (n-1)^2 C_{n-2} \text{ with } r_0 = 1 \text{ and } r_1 = 2$$

According to [12] the sequences 1, 4, 22, 149, 1186, ... given the formular

$$|A_n^c| = \frac{n!}{2} + \frac{n^2}{2} (n-1)! + \sum_{r=0}^{n-2} \binom{n}{r}^2 r!$$

For $n=1, 2, 3, \dots$

Main Result

Identification of Idempotent and Nilpotent Elements in Alternating Semigroup

In this paper, we show that $|EA_n^e| = 2^n$ for $n \geq 2$.

Where $|EA_n^e| = 1$

And

$$|NA_n^e| = \begin{cases} \sum_{r=0}^{n-1} \binom{n}{r} \binom{n-1}{r} r! & \text{if } n \text{ is odd} \\ \sum_{r=0}^{n-2} \binom{n}{r} \binom{n-1}{r} r! & \text{if } n \text{ is even} \end{cases}$$

Where $|EA_n^e|$ and $|NA_n^e|$ are the number of idempotent and nilpotent elements in alternating semigroup respectively.

The idempotent and nilpotent elements in A_n^e are identified and marked (-) and (*) respectively.

The List

The elements in A_1^e

For $n=1$, has 1 element.

A_1^e	{1}
	$\left(\begin{smallmatrix} 1^{*-} \\ - \end{smallmatrix} \right)$

The elements in A_2^e

For $n=2$, has 4 elements

A_2^e	{1, 2}
	$\left(\begin{smallmatrix} 12^- \\ 12 \end{smallmatrix} \right) \left(\begin{smallmatrix} 12^- \\ 1- \end{smallmatrix} \right) \left(\begin{smallmatrix} 12^- \\ -2 \end{smallmatrix} \right) \left(\begin{smallmatrix} 12^- \\ -- \end{smallmatrix} \right)$

The elements in A_3^e

for $n=3$, has 22 elements.

A_3^e	{1, 2, 3}
	$\left(\begin{smallmatrix} 123^- \\ 123 \end{smallmatrix} \right) \left(\begin{smallmatrix} 123 \\ 231 \end{smallmatrix} \right) \left(\begin{smallmatrix} 123 \\ 312 \end{smallmatrix} \right) \left(\begin{smallmatrix} 123^- \\ 12- \end{smallmatrix} \right) \left(\begin{smallmatrix} 123^* \\ -12 \end{smallmatrix} \right) \left(\begin{smallmatrix} 123^- \\ 1-3 \end{smallmatrix} \right) \left(\begin{smallmatrix} 123^* \\ 2-1 \end{smallmatrix} \right) \left(\begin{smallmatrix} 123^* \\ 23- \end{smallmatrix} \right)$
	$\left(\begin{smallmatrix} 123^- \\ -23 \end{smallmatrix} \right) \left(\begin{smallmatrix} 123^* \\ 31- \end{smallmatrix} \right) \left(\begin{smallmatrix} 123^* \\ -31 \end{smallmatrix} \right) \left(\begin{smallmatrix} 123^* \\ 3-2 \end{smallmatrix} \right) \left(\begin{smallmatrix} 123^- \\ 1-- \end{smallmatrix} \right) \left(\begin{smallmatrix} 123^* \\ -1- \end{smallmatrix} \right) \left(\begin{smallmatrix} 123^* \\ --1 \end{smallmatrix} \right)$
	$\left(\begin{smallmatrix} 123^* \\ 2-- \end{smallmatrix} \right) \left(\begin{smallmatrix} 123^- \\ -2- \end{smallmatrix} \right) \left(\begin{smallmatrix} 123^* \\ --2 \end{smallmatrix} \right) \left(\begin{smallmatrix} 123^* \\ 3-- \end{smallmatrix} \right) \left(\begin{smallmatrix} 123^* \\ -3- \end{smallmatrix} \right) \left(\begin{smallmatrix} 123^- \\ --3 \end{smallmatrix} \right) \left(\begin{smallmatrix} 123^* \\ --- \end{smallmatrix} \right)$

The elements of alternating semigroup of A_4^e and A_5^e were also generated but because of the space we could not display it. The table below shows our results

Table 2

n	$ A_n^e $	$ EA_n^e $	$ NA_n^e $
1	1	1	1
2	4	4	1
3	22	8	13
4	149	16	49
5	1186	32	501

From the sequences 1, 4, 8, 16, 32,... and 1, 1, 13, 49, 501,... in table 2, we were able to derived the formular for the numbers of Idempotent and Nilpotent elements in Alternating semigroups as

$$|EA_n^e| = 2^n \text{ for } n \geq 2.$$

and

$$|NA_n^e| = \begin{cases} \sum_{r=0}^{n-1} \binom{n}{r} \binom{n-1}{r} r! & \text{if } n \text{ is odd} \\ \sum_{r=0}^{n-2} \binom{n}{r} \binom{n-1}{r} r! & \text{if } n \text{ is even} \end{cases}$$

Discussion and Conclusion

In this paper, partial one-one transformation semigroups has been successfully used to identified the number of elements of alternating semigroup $|A_n^e|$ and the formular for the number of elements of idempotent and nilpotent were derived. The proof for the formular obtained is still in progress. Based on this paper, we have strong believe that many things can be deduced if further studies could be done. The theory of semigroup has it's scope widened to embrace many aspects of theoretical computer sciences, such as: *automata theory, coding theory, computational theory and formal languages* as well as applications in the sciences. It's also assist people and (Computers) in sorting data and designing better networks.

Acknowledgement: we would like to gratefully acknowledge the support of Dr. A. Umar, Department of Mathematics and Statistics Qaboos University OMAN.

References

- Borwein, (1989). The number of semigroup in partial one-one transformation.
- Borwein, (1989). The number of semigroup of order decreasing partial one-one transformation.
- Borwein, (1989). The number of Idempotent element $|E(S)|$ of the semigroup of order decreasing partial one-one transformation.
- Garba, G. U., (1994). The number of semigroup of the order-preserving partial one-one Transformation.
- Garba, G. U. (1994). The number of Idempotent element $|E(S)|$ for the order preserving partial one-one transformation semigroup.
- The Ganyuushkin (2003). The number of semigroup of the order-preserving and order-decreasing partial one-one transformation.
- The Ganyuushkin (2003). The number of semigroup of Idempotent element $|E(S)|$ of the order preserving and order-decreasing partial one-one transformation semigroup.
- Laradji (2007). The number of nilpotent element $|N(S)|$ in partial one-one transformation
- Laradji, A, & Umar, A. (2007). Combinatorial results for the symmetric inverse semigroup. Semigroup forum, 75, 221-236.
- Umar, A., (1992). It on the number of nilpotent element $|N(S)|$ of the order-decreasing partial one-one transformation.

Umar, A. (2004). It on the number of idempotent element $|E(S)|$ in partial one-one transformation

Stephen, L. (1996). It Symmetric inverse semigroups. *Mathematical survey and monography*.